

STUDIES IN STRUCTURE AND PROPERTIES
OF LIGHT NUCLEI

by

Santosh K. Shah

For the degree of

DOCTOR OF PHILOSOPHY

of

THE M.S. UNIVERSITY OF GUJARAT

043



B2044

December 1962

Physical Research Laboratory

Ahmedabad - 9, INDIA

PREFACE

A major problem in the theory of nuclei is to obtain an understanding of the nuclear forces operating within nuclei as distinct from the forces between two free nucleons. Despite the large amount of work by Brueckner and many others (which has helped to clarify and, to a large extent, justify the conceptual basis of nuclear shell model), calculations of nuclear properties, such as energy levels starting with free nucleon-nucleon interactions, present considerable mathematical complexity and are not in general easy to carry out without the aid of large computers. As a result, most shell model calculations are in practice done by assuming a relatively simple effective interaction. In this dissertation an attempt is made to study in a systematic way the nature of this effective interaction in several simple nuclei.

The work of Brueckner and others shows that the effective nuclear forces are non-local or coordinate-dependent and also configuration-dependent. However, they are non-singular, i.e., will not have an infinite repulsive core, as is generally assumed in free nucleon forces. With these considerations in mind, we suggest in chapter II a method by which nuclear spectra may be calculated, and which may exhibit clearly the above properties of the effective forces. The idea is to

express the matrix elements of the nuclear hamiltonian in terms of the matrix elements of the interaction in two-particle states of different spin, isotopic spin and relative orbital angular momentum. This procedure would be somewhat analogous to the analysis of the nucleon-nucleon scattering data in terms of phase shifts in different states of the nucleons. It is pointed out that, if enough experimental data on energy levels is available, it should be possible to study directly the nature of the effective interaction in different spin and 1-spin states and, in particular, a single type of momentum-dependence, via., angular-momentum-dependence of the forces, can be easily brought out. The formalism for such calculations is described in chapter III, and is then applied to some simple nuclei such as Li^{10} , 6_2He and $^2 = 1$ states in Li^7 . Although the experimental data is not sufficient at present to make more definite analysis, it is pointed out that the even-state (singlet) interactions appear to be relatively configuration-independent, whereas the odd-state (triplet) forces appear to vary from nuclei to nuclei. More data will not only enable us to make these statements more precise, but will also enable us to extend the analysis to other states (such as $1 = 0$ states), and other heavier nuclei. We would like to stress the importance of our method of analyzing the effective interaction in states of relative orbital angular momentum which should be more powerful and useful to

illustrate the properties of the effective interaction rather than the conventional method of expressing the interaction in terms of existing multipole.

In chapter III some additional considerations are presented for the special case of isotopes O^{18} and O^{19} . In the first section, the results of our analysis of chapter II are compared with the results of other authors who made calculations of O^{18} spectrum. Next, the parameters of the effective interaction are tested by calculating the energy levels of O^{19} . It is found in chapter II that several different sets of parameters could give equally satisfactory results for O^{18} . Here we find that, if some of the excited states of O^{19} have their spins and parities measured, it may be possible to distinguish between some of these different sets of parameters. Finally, in section 3, the available data on the energy levels of O^{18} and O^{19} (eight levels) is re-analysed (by partial inversion of the energy matrices) to determine the matrix elements of the interaction in O^{18} states. This provides an additional check on the interaction parameters discussed in chapter II.

At the end of the thesis, reprints of our published work are included. These refer to the work included in the thesis as well as some work (nuclear interactions in $3/2^-$ -doublets, etc.) not included in the thesis in the interest of homogeneity.

ACKNOWLEDGEMENT

The author is indebted to Dr. S. Pandya for introducing him to the subject, for his guidance and helpful discussions during the progress of the work. He would also like to record his indebtedness to his colleagues Mr. V. K. Shankappan, Mr. Y. R. Vaghmare and Mr. V. L. Maralikar for their kind cooperation and useful discussions.

He would like to express his gratitude to Professor T. R. Narayanan for having extended to him the hospitality of the Physical Research Laboratory. His thanks are also due to Professor H. G. Pandya for his encouragement and interest during the course of this work.

Finally, it is a pleasure to acknowledge the award of Research Scholarship by the M.S. University of Baroda, Baroda, during the work for the thesis.

CONTENTS

preface	(1)
Acknowledgements	(iv)
CHAPTER I. Introduction.	
1. Nuclear Shell Model	1
2. Brueckner Theory	9
2a. Realistic Two-Body Potential	9
2b. Reaction Matrix	12
3. Scope of the Thesis	18
CHAPTER II. Effective Interactions in Nuclei.	
1. Introduction	21
2. Method of Analysis	24
3. Pairing Energies	33
4. Analysis of closed shell + 2 Nucleon Configurations	37
5. Conclusions	54
CHAPTER III. An Analysis of the Energy Levels of α^{18} and α^{19} .	
1. Introduction	57
2. Comparison with results of other authors	58
3. Energy Levels of α^{19} (I)	68
4. Energy Levels of α^{19} (II)	74
CHAPTER IV. Concluding Remarks	
References	80
Appendix - Reprints of the published papers.	88